

# Graphs with Maximum Edge-Integrity

Lowell W. Beineke, Indiana-Purdue University at Fort Wayne

Wayne Goddard, University of Pennsylvania

Marc J. Lipman, Office of Naval Research

## Abstract

The edge-integrity of a graph  $G$  is given by the minimum of  $|S| + m(G - S)$  taken over all  $S \subseteq E(G)$ , where  $m(G - S)$  denotes the maximum order of a component of  $G - S$ . An honest graph is one with maximum edge-integrity (viz. its order). In this paper lower and upper bounds on the edge-integrity of a graph with given order and diameter are investigated. For example, it is shown that the diameter of an honest graph on  $n$  vertices is at most  $\sqrt{8n} - 3$ , and this is sharp. Also, a lower bound for the edge-integrity of a graph in terms of its eigenvalues is established. This is used to show that for  $d$  sufficiently large almost all  $d$ -regular graphs are honest.

Correspondence to:  
Wayne Goddard  
Dept of Mathematics  
University of Pennsylvania  
Philadelphia PA 19103  
wgoddard@math.upenn.edu

## 1 Introduction

In this paper we consider finite undirected graphs without loops or multiple edges. The edge-integrity of a graph attempts to measure the disruption caused by the removal of edges from the graph. The *order* of a component or graph is the number of its vertices, and we let  $m(H)$  denote the maximum order of a component of graph  $H$ . Barefoot, Entringer and Swart [3] defined the *edge-integrity* of a graph  $G$  with edge set  $E(G)$  by

$$I'(G) = \min_{S \subseteq E(G)} |S| + m(G - S).$$

Any set  $S$  of edges which realizes this value is called an  *$I'$ -set*; one of minimum cardinality is called a *minimum  $I'$ -set*. We say that a graph is *honest* if its edge-integrity is equal to its order. Of course, the empty set shows that  $I'(G) \leq m(G)$  in general.

Barefoot, Entringer and Swart [3] proposed this parameter as a measure of how hard it is to disrupt thoroughly a network by edge failures. Properties of edge-integrity were also investigated by Bagga et al., and by others. See for example [4, 6, 8, 10]. There is also a survey [5].

In this paper we investigate the range of values that the edge-integrity may take given the order and diameter of a graph. One of the first results in this direction was given by Bagga et al. who showed that graphs with diameter 2 are honest:

**Proposition 1** [4] *If  $G$  has  $n$  vertices and diameter 2 then  $I'(G) = n$ .*

For graphs with diameter 3 we prove a sharp lower bound of  $3n^{2/3}/2 - O(n^{1/3})$  on the edge-integrity of such graphs. But, for graphs with higher diameter there is no better lower bound than that was observed by Barefoot et al.:

**Proposition 2** [3] *If  $G$  has  $n$  vertices and is connected then  $I'(G) \geq \lceil 2\sqrt{n} \rceil - 1$ .*

They showed that the path  $P_n$  on  $n$  vertices has  $I'(P_n) = \lceil 2\sqrt{n} \rceil - 1$ . We show that there are graphs with diameter 4 (and indeed with radius 2) with the same edge-integrity. At the other extreme, we show the diameter of an honest graph on  $n$  vertices is at most  $\sqrt{8n} - 3$ , and this is sharp.

In the final section we establish a link between the edge-integrity of a graph and its eigenvalues. As a consequence we show that, for  $d$  sufficiently large, almost all  $d$ -regular graphs are honest.

## 2 Minimum Edge-Integrity and Diameter

We know already by Proposition 1 that a graph with diameter 2 is honest. Our first result gives a tight lower bound on the edge-integrity of a graph of diameter 3:

**Theorem 1** *Let graph  $G$  have  $n$  vertices and diameter 3. Then*

$$I'(G) \geq 3n^{2/3}/2 - n^{1/3}/2 - O(1).$$

PROOF. Let  $S$  be an  $I'$ -set of  $G$ . If every vertex of  $G$  is incident with an edge of  $S$  then  $|S| \geq n/2$  and we are done. Otherwise, let  $H_1, H_2, \dots, H_t$  be the components of  $G - S$  which contain a vertex that is not incident to an edge of  $S$ . Then, as the diameter of  $G$  is at most 3,  $S$  contains an edge between  $H_i$  and  $H_j$  for  $1 \leq i < j \leq t$ . Let  $H_1, H_2, \dots, H_t$  have a total of  $r$  vertices. Then  $|S| \geq \binom{t}{2} + (n - r)/2$ . And  $m(G - S) \geq r/t$ .

Thus  $I'(G)$  is at least the minimum of

$$\frac{r}{t} + \binom{t}{2} + (n - r)/2,$$

taken over  $1 \leq t \leq r \leq n$ . For  $t = 1$  the minimum of the expression is  $(n + 1)/2$ . For  $t \geq 2$  the minimum can be determined by using calculus (and a computer): it is attained at  $r^* = n$  and  $t^* \approx n^{1/3}$ , and has the above value. QED

There are graphs of diameter 3 which have edge-integrity that matches the lower bound. For example, for  $t$  even let  $G_t$  be the graph formed by taking  $t$  disjoint cliques, each with  $t^2 - t/2$  vertices, and adding one edge between every pair of cliques. The graph  $G_t$  has  $n_t = t^3 - t^2/2$  vertices and edge-integrity  $i_t = 3t^2/2 - t$ . The limit of  $i_t - (3n_t^{2/3}/2 - n_t^{1/3}/2)$  as  $t$  goes to infinity is  $-1/24$ .

If the diameter is 4, however, then it turns out that the edge-integrity can be as small as what connectivity guarantees (recall Proposition 2). For example, construct graph  $H_s$  as follows. Take  $s$  disjoint cliques, each with  $s$  vertices, and designate one vertex in each clique; then add  $s - 1$  edges between the designated vertices to form a star. The resulting graph  $H_s$  has radius 2 and the same edge-integrity as the path on  $s^2$  vertices, viz.  $2s - 1$ . The graph  $H_4$  is illustrated in Figure 1.

## 3 Maximum Edge-Integrity and Diameter

For graphs with large edge-integrity and large diameter consider the following graphs. Given a sequence  $a_0, a_1, \dots, a_d$  of positive integers, we define the ‘‘sausage’’




Figure 1: The graph  $H_4$  has radius two and minimum edge-integrity

graph  $G[a_0, a_1, \dots, a_d]$  as follows: take disjoint cliques  $A_0, A_1, \dots, A_n$  where  $A_i$  has  $a_i$  vertices ( $i = 0, 1, \dots, d$ ), and add all edges between  $A_i$  and  $A_{i+1}$  for  $i = 0, 1, \dots, d - 1$ . Figure 2 shows  $G[1, 1, 2, 2, 2, 1, 1]$ . Of course,  $G[a_0, a_1, \dots, a_d]$  has diameter  $d$ .




Figure 2: An honest sausage graph of diameter 6

The following lemma aids in the calculation of the edge-integrity of sausage graphs:

**Lemma 1** *Let  $G = G[a_0, a_1, \dots, a_d]$  be a sausage graph, and let  $S$  be a minimum  $I'$ -set of  $G$ . Then*

- (a) *the removal of  $S$  does not split any of the  $A_i$ , and*
- (b) *if the removal of  $S$  separates  $A_i$  and  $A_{i+1}$  then  $a_i \leq a_{i+2}$  and  $a_{i+1} \leq a_{i-1}$ .*

PROOF. (a) Suppose the removal of  $S$  splits  $A_i$ . Let  $H_1$  and  $H_2$  be two components of  $G - S$  that contain vertices of  $A_i$ . Let  $B_r = \{j : A_j \cap H_r \neq \emptyset\}$  for  $r = 1, 2$ . Two cases arise:

1. *One of the  $B_r$  is a subset of the other. Say  $B_1 \subseteq B_2$ . Then combine  $H_1$  and  $H_2$ ; that is, expunge from  $S$  and add to  $G - S$  all the edges that join vertices of  $H_1$  and  $H_2$ . The number of edges expunged from  $S$  is at least  $|H_1|$ , while the increase in the maximum component order is at most  $|H_1|$ . Thus  $S$  was not a minimum  $I'$ -set, a contradiction.*

2. Neither of the  $B_r$  is a subset of the other. Say  $i-1 \in B_1 - B_2$  and  $i \in B_1 \cap B_2$ . Let  $T = H_1 \cap A_{i-1}$ ,  $U = H_1 \cap A_i$ ,  $V = H_2 \cap A_i$ , and  $W = H_2 \cap A_{i+1}$ . If  $|W| \leq |T|$  then add to  $G - S$  the edges between  $V$  and  $T \cup U$  and remove the edges between  $V$  and  $W$ . Effectively this transfers  $V$  from  $H_2$  to  $H_1$ , while saving at least  $|V|$  edges. Thus  $S$  was not a minimum  $I'$ -set.

If  $|W| > |T|$  then remove from  $G - S$  the edges between  $U$  and  $T$ , splitting  $H_1$  into two pieces, and add all edges between the piece of  $H_1$  containing  $U$  and  $H_2$ . It can be checked that the net decrease in the number of edges removed is again at least the increase in the maximum component order. Thus  $S$  was not a minimum  $I'$ -set.

(b) Now suppose  $S$  separates  $A_i$  and  $A_{i+1}$ . If  $a_{i-1} < a_{i+1}$  then we can move the cut to separate  $A_i$  and  $A_{i-1}$  rather. The saving in edges removed is at least  $|A_i|$ , while the increase in maximum component order is at most  $|A_i|$ . So  $S$  was not a minimum  $I'$  set. QED

**Corollary 2** *Let  $G[a_0, a_1, \dots, a_d]$  be a sausage graph for which there exists an  $r$  such that for  $0 \leq i \leq r - 2$  it holds that  $a_i < a_{i+2}$ , and for  $r \leq i \leq d - 2$  it holds that  $a_i > a_{i+2}$ . Then  $G$  is honest.*

PROOF. By the above lemma a minimum  $I'$ -set of the graph is empty. QED

For example the sausage graph  $G[1, 1, 2, 2, 2, 1, 1]$  of Figure 2 is honest (use  $r = 3$ ). We will show that the cheapest way to satisfy the hypothesis of the corollary gives the honest graph of diameter  $d$  with minimum order. We will need the following lemma:

**Lemma 2** *Let  $m$  be a given integer, and consider the following problem:*

$$\text{Minimize } \sum_{i=0}^m b_i \text{ such that } b_j b_{j+1} \geq \sum_{i=0}^j b_i \text{ for } j = 0, 1, \dots, m-1,$$

where the  $b_i$  are positive integers. Then the minimum is  $\lfloor (m+2)^2/4 \rfloor$ , and the unique best sequence of the  $b_i$  is given by  $\mathcal{B} = 1, 1, 2, 2, 3, 3, 4, 4, \dots$

PROOF. Let  $\beta_j = \sum_{i=0}^j b_i$ . Rearranged, the constraint says that  $b_j(b_{j+1}-1) \geq \beta_{j-1}$ . So  $b_j + (b_{j+1}-1) \geq \lceil 2\sqrt{\beta_{j-1}} \rceil$ . Thus  $\beta_{j+1} = b_{j+1} + b_j + \beta_{j-1} \geq 1 + \lceil 2\sqrt{\beta_{j-1}} \rceil + \beta_{j-1}$ . By induction it then follows that  $\beta_j \geq \lfloor (j+2)^2/4 \rfloor$ . (We omit the straight-forward calculation.) QED

**Theorem 3** *The minimum number of vertices in an honest graph of diameter  $d$  is  $\lceil (d+2)(d+4)/8 \rceil$ .*

PROOF. Let  $G$  be an honest graph with  $n$  vertices and diameter  $d$ . Let  $v$  be a vertex such that there is a vertex at distance  $d$  from  $v$ . Let  $A_i$  denote the set of vertices at distance  $i$  from  $v$ , and let  $a_i = |A_i|$ . Further let  $r$  be such that  $\sum_{i=1}^r a_i \leq n/2$  and  $\sum_{i=r+1}^d a_i \leq n/2$ .

By considering the removal of all edges between  $A_j$  and  $A_{j+1}$ , it follows that necessarily

$$a_j a_{j+1} \geq \sum_{i=0}^j a_i \quad \text{for } j = 1, \dots, r,$$

and

$$a_j a_{j+1} \geq \sum_{i=j+1}^d a_i \quad \text{for } j = r, \dots, d-1.$$

By the above lemma it follows that  $\sum_{i=0}^r a_i \geq \lfloor (r+2)^2/4 \rfloor$ , and  $\sum_{i=r+1}^d a_i \geq \lfloor (d-r+1)^2/4 \rfloor$ . So it follows that

$$8n \geq 2(r+2)^2 + 2(d-r+1)^2 - 8\varepsilon_{dr},$$

where  $\varepsilon_{dr} = 0$  if  $d, r$  both even,  $\varepsilon_{dr} = 1/2$  if  $d, r$  both odd, and  $\varepsilon_{dr} = 1/4$  otherwise.

If  $d$  is even, then the lower bound for  $8n$  is minimized at  $r = d/2$  or  $r = d/2 - 1$ , depending on which is odd. It follows that  $8n \geq d^2 + 6d + 8$ , as required. If  $d$  is odd the lower bound for  $8n$  ignoring the  $\varepsilon_{dr}$ -term is minimized at  $r = (d-1)/2$  only where it has value  $8n \geq d^2 + 6d + 9$ .

It can be checked that the only way the RHS expression can go below  $d^2 + 6d + 8$  is if  $d$  is odd and  $r = (d-1)/2$  is odd. But by the above lemma it also follows that  $\sum_{i=0}^{r-1} a_i \geq \lfloor (r+1)^2/4 \rfloor$  and  $\sum_{i=r}^d a_i \geq \lfloor (d-r+2)^2/4 \rfloor$ . From this it can be shown that the desired lower bound is valid.

To obtain a best sequence of the  $a_i$  we put together two almost equal-sized initial segments of  $\mathcal{B}$  with the second one reversed. If  $d$  is even the two initial segments differ in length by 1. If  $d \equiv 1 \pmod{4}$  then the two initial segments have the same length. If  $d \equiv 3 \pmod{4}$  then the two initial segments differ in length by 2. For example, for  $d = 6$  the best  $\{a_i\}$  is 1, 1, 2, 2, 2, 1, 1. For  $d = 7$  it is 1, 1, 2, 2, 3, 2, 1, 1. The associated sausage graphs are honest by Corollary 2. QED

We believe that ‘‘honest sausage graphs with tails’’ have nearly maximum edge-integrity for their order and diameter. Let

$$G_d^t = G[\underbrace{1, 1, \dots, 1}_t, \underbrace{1, 1, 2, 2, 3, \dots, 3, 2, 2, 1, 1, 1, \dots, 1}_d, \underbrace{1, 1, \dots, 1}_t],$$

where the middle portion gives the honest graph  $G_d$  of minimum order  $n_d$  described in Theorem 3. Then  $G_d^t$  has diameter  $D = 2t + d$  and order  $N = n_d + 2t$ . By

Lemma 1, it follows that the subgraph  $G_d$  remains virtually intact after the removal of a minimum  $I'$ -set  $S$  (except maybe losing one vertex at both ends). So  $m(G_d^t - S) \geq n_d - 2$ . Calculations then show that

$$I'(G_d^t) = \begin{cases} n_d, & n_d \geq t + 3; \\ \approx n_d - 2 + (2t + 2)/(n_d - 2), & \sqrt{2t} \leq n_d \leq t + 3; \\ \approx I'(P_{D+1}), & n_d \leq \sqrt{2t}. \end{cases}$$

Specifically, we conjecture:

**Conjecture 1** *For  $p_d \geq t + 3$ , the graph  $G_d^t$  has the maximum edge-integrity for a graph of its order  $N$  and diameter  $D$ .*

That edge-integrity is approximately  $(N - D) + \sqrt{8(N - D)} + 1$ , and the range of appropriate  $D$  is roughly  $\sqrt{8N} - 3 \leq D \leq 2N/3$ .

## 4 Edge-Integrity and Eigenvalues

In this section we derive a simple lower bound on the edge-integrity of a graph in terms of its eigenvalues.

Let  $G$  be a graph and  $A$  a subset of the vertices. Then the (*edge*) *boundary*  $b(A)$  of  $A$  is the number of edges of  $G$  with exactly one end in  $A$ . We let  $b(m)$  denote the minimum of  $b(A)$  taken over sets  $A$  of  $m$  vertices. Isoperimetric inequalities give lower bounds for edge-integrity, as was observed in [7]:

**Proposition 3** [7] *Let  $G$  be a graph on  $n$  vertices and let  $f(x)$  be a real convex function such that  $f(m) \leq b(m)$  for all  $m \in \{1, 2, \dots, n\}$ . Then*

$$I'(G) \geq \min_{x \geq 0} x + \frac{n}{2x} f(x).$$

Alon and Milman [1] established a link between the boundary and eigenvalues. Let  $L$  denote the Laplacian matrix  $D - A$  of the graph, where  $A$  is the adjacency matrix of the graph, and  $D$  a diagonal matrix with the degrees of the vertices on the diagonal. Then the eigenvalues of  $L$  are real and nonnegative. Let  $\lambda_1$  denote the second smallest eigenvalue. (The smallest is 0.) Alon and Milman showed:

**Proposition 4** [1] *For a graph on  $n$  vertices it holds that  $b(m) \geq \lambda_1 m(1 - m/n)$ .*

A corollary of the above two results is:

**Theorem 4** For a graph  $G$  on  $n$  vertices whose Laplacian has second smallest eigenvalue  $\lambda_1$ ,

$$I'(G) \geq n \cdot \min(1, \lambda_1/2).$$

PROOF. By the above two propositions,

$$I'(G) \geq \min_x x + n \frac{\lambda_1 x(1 - x/n)}{2x},$$

where the minimum is taken over real  $x \in [0, n]$ . The minimum is attained either at  $x = n$ , where it has value  $n$ , or at  $x = 0$ , where it has value  $n\lambda_1/2$ . QED

For example, all the hypercubes have  $\lambda_1 = 2$  and are thus honest. This was first shown by Bagga et al. [4].

In [11] the third author showed that there are only finitely many cubic graphs which are honest. In contrast, when  $d$  is sufficiently large, almost every  $d$ -regular graph is honest. For, Friedman [9] showed that for a random  $d$ -regular graph almost surely  $\lambda_1 \geq d - 2\sqrt{d-1} - O(\log d)$ . And hence for  $d$  sufficiently large, by Theorem 4 the graph is almost surely honest.

## Acknowledgements

The research of the first author was supported in part by ONR contract N00014-86-K-0412 and the second author by ONR contract N00014-91-J-1022. The second author would also like to thank IPFW for its hospitality, DIMACS for financial support, and Nabil Kahale for helpful discussions.

## References

- [1] N. ALON & V.D. MILMAN,  $\lambda_1$ , isoperimetric inequalities for graphs, and superconcentrators, *J. Combin. Theory Ser. B* **38** (1985), 73–88.
- [2] C.A. BAREFOOT, R. ENTRINGER & H.C. SWART, Vulnerability in graphs—A comparative survey, *J. Combin. Math. Combin. Comput.* **1** (1987), 13–22.
- [3] C.A. BAREFOOT, R. ENTRINGER & H.C. SWART, Integrity of trees and powers of cycles, *Congress. Numer.* **58** (1987), 103–114.
- [4] K.S. BAGGA, L.W. BEINEKE, M.J. LIPMAN & R.E. PIPPERT, On the edge-integrity of graphs, *Congress. Numer.* **60** (1987), 141–144.
- [5] K.S. BAGGA, L.W. BEINEKE, M.J. LIPMAN & R.E. PIPPERT, Edge-integrity: a survey, *Discrete Math.* **124** (1994), 3–12.

- [6] K.S. BAGGA, L.W. BEINEKE, M.J. LIPMAN, R.E. PIPPERT & R.L. SEDLMAYER, A good algorithm for the computation of the edge-integrity of trees, *Congress. Numer.* **67** (1988), 225–232.
- [7] L.W. BEINEKE, W. GODDARD & M.J. LIPMAN, The edge-integrity of cartesian products, submitted.
- [8] M.R. FELLOWS & S. STUECKLE, The immersion order, forbidden subgraphs and the complexity of network integrity, *J. Combin. Math. Combin. Comput.* **6** (1989), 23–32.
- [9] J. FRIEDMAN, On the second eigenvalue and random walks in random  $d$ -regular graphs, *Combinatorica* **11** (1991), 331–362.
- [10] R. LASKAR, S. STUECKLE & B. PIAZZA, On the edge-integrity of some graphs and their complements, *Discrete Math.* **122** (1993), 245–253.
- [11] M.J. LIPMAN, The honest cubic graphs, *Congress. Numer.* **85** (1991), 184–192.